

## "Long-term non-surgical control with ulipristal acetate of multiple uterine fibroids, enabling pregnancy."

Luyckx, Mathieu ; Pirard, Céline ; Fellah, Latifa ; Dereume, Alice ;  
Mhallem Gziri, Mina ; Debiève, Frédéric ; Squifflet, Jean-Luc

Document type : *Article de périodique (Journal article)*

---

### Référence bibliographique

Luyckx, Mathieu ; Pirard, Céline ; Fellah, Latifa ; Dereume, Alice ; Mhallem Gziri, Mina ; et. al. *Long-term non-surgical control with ulipristal acetate of multiple uterine fibroids, enabling pregnancy..* In: *American Journal of Obstetrics and Gynecology*, Vol. 214, no. 6, p. 756.e1-756.e2 (2016)

DOI : 10.1016/j.ajog.2016.02.049

# Accepted Manuscript

Long-term non-surgical control with ulipristal acetate of multiple uterine fibroids, enabling pregnancy

Dr. Mathieu Luyckx, MD, Prof Céline Pirard, PhD, Dr Latifa Fellah, MD, Dr Alice Dereume, MD, Prof Mina Mhallem, PhD, Prof Frédéric Debiève, PhD, Prof Jean Squifflet, PhD

PII: S0002-9378(16)00391-4

DOI: [10.1016/j.ajog.2016.02.049](https://doi.org/10.1016/j.ajog.2016.02.049)

Reference: YMOB 10973

To appear in: *American Journal of Obstetrics and Gynecology*

Received Date: 9 February 2016

Revised Date: 22 February 2016

Accepted Date: 24 February 2016

Please cite this article as: Luyckx M, Pirard C, Fellah L, Dereume A, Mhallem M, Debiève F, Squifflet J, Long-term non-surgical control with ulipristal acetate of multiple uterine fibroids, enabling pregnancy, *American Journal of Obstetrics and Gynecology* (2016), doi: 10.1016/j.ajog.2016.02.049.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



# Long-term non-surgical control with ulipristal acetate of multiple uterine fibroids, enabling pregnancy

**Dr. Mathieu Luyckx, MD:** - Gynecology and Andrology Department, Cliniques Universitaires Saint-Luc, Brussels, Belgium

- Institut de Recherche Expérimentale et Clinique (IREC),  
Université Catholique de Louvain, Brussels, Belgium

**Prof Céline Pirard PhD:** - Gynecology and Andrology Department, Cliniques Universitaires Saint-Luc, Brussels, Belgium

- Institut de Recherche Expérimentale et Clinique (IREC),  
Université Catholique de Louvain, Brussels, Belgium

**Dr Latifa Fellah MD:** Radiology Department, Cliniques Universitaires Saint-Luc, Brussels, Belgium

**Dr Alice Dereume MD:** - Gynecology and Andrology Department, Cliniques Universitaires Saint-Luc, Brussels, Belgium

**Prof Mina Mhallem, PhD:** - Obstetrics Department, Cliniques Universitaires Saint-Luc, Brussels, Belgium

- Institut de Recherche Expérimentale et Clinique (IREC),  
Université Catholique de Louvain, Brussels, Belgium

**Prof Frédéric Debiève PhD:** - Obstetrics Department, Cliniques Universitaires Saint-Luc, Brussels, Belgium

- Institut de Recherche Expérimentale et Clinique (IREC),  
Université Catholique de Louvain, Brussels, Belgium

**Prof Jean Squifflet, PhD:** - Gynecology and Andrology Department, Cliniques Universitaires Saint-Luc, Brussels, Belgium

- Institut de Recherche Expérimentale et Clinique (IREC),  
Université Catholique de Louvain, Brussels, Belgium

Declaration of interests: None of the authors have any interests to declare.

Corresponding author: Mathieu Luyckx MD,

Gynecology and Andrology Department, Cliniques Universitaires Saint-Luc,

Avenue Hippocrate 10, B-1200 Brussels, Belgium;

Tel: +32 2 764 95 09

**mathieu.luyckx@uclouvain.be**

Condensation:

MR images depict a marked response to Ulipristale Acetate in a woman with large uterine fibroids, and demonstrate no significant regrowth during pregnancy or 12 weeks postpartum.

Short Title: UPA non-surgical fibroids control

Case note: A 39-year-old woman under our care for five years for multiple uterine fibroids presented requesting a review of her condition with a view to attempting pregnancy. Previous treatment included laparotomic myomectomy in 2006, GnRH agonist therapy (monthly leuprolide acetate injections in the PEARL II study<sup>1</sup>) in 2008, and hysteroscopic myomectomy in 2009, and had been asymptomatic on desogestrel treatment for 1 year. On clinical examination, uterine volume was equivalent to 20 weeks' gestation and magnetic resonance imaging (MRI) revealed multiple, disseminated, diffuse myomatosis (Figure A). Treatment was initiated with 5 mg oral ulipristal acetate (UPA) daily. MRI evaluation showed a significant decrease in fibroid volume after 3 months, continuing after 6 months (Figures B, C). Without surgery, the patient proceeded with in vitro fertilization for andrologic concern and age. Pregnancy was confirmed following her first embryo transfer, 11 months after beginning and 3.5 months after ending UPA treatment. No significant fibroid regrowth was noted at MRI performed during pregnancy (Figures D, E). Because of previous laparotomy with myomectomy, cesarean section was planned and a healthy baby was delivered at 39 weeks. Follow-up MRI 12 weeks after delivery showed sustained regression of the myomas, both in number and size, also due to 'natural' remodeling after pregnancy<sup>2</sup> (Figure F). The patient remains asymptomatic one year post-delivery.

Comment: This MRI-based analysis of UPA treatment for diffuse uterine myomatosis reveals an unprecedented level of effectiveness, sustained even throughout the subsequent pregnancy. However, fibroid response to UPA in terms of volume reduction is very unpredictable<sup>3</sup>. Courtoy et al. conducted immunohistochemistry, microarrays and histology studies on surgical specimens and described a multifactorial mechanism of action<sup>4</sup>: UPA increased apoptosis of myoma cells, but only at the beginning of treatment, by a pathway independent of the caspase. After long-term treatment with UPA, modification of the extra-cellular matrix (ECM) was significant compared to the untreated group, with an increase in matrix

metalloproteinase 2 (MMP-2) levels. Hence, in patients showing such a marked response to UPA treatment, it is possible that pregnancy could continue to modify the ECM by uterine growth. The remaining fibroid cells may have been selected by primary UPA-mediated apoptosis resulting in a population of cells less sensitive to hormonal stimulation, due to modification of progesterone receptor expression, as shown for other selective progesterone receptor modulators<sup>5</sup>.

Acknowledgements : The authors thank Laurence Beausaert, research nurse in charge of the UPA studies and Mira Hryniuk for reviewing the English language of the manuscript

**Figure:** *MRI images showing regression of uterine fibroids during UPA treatment and subsequent pregnancy*

Multiple large myomas (dark ovals) and a distorted and reduced-capacity uterine cavity (light area) were visible before treatment (A). Subsequent MRIs showed successive improvements after 3 months (B) and 6 months (C) of UPA; with no significant regrowth during early pregnancy (D, 13 weeks' gestation). Further shrinkage was observed in the third trimester (E, 32 weeks' gestation), with disappearance of most fibroids by 12 weeks after delivery (F).

## REFERENCES

- <sup>1</sup> Donnez J, Tomaszewski J, Vázquez F, et al. Ulipristal acetate versus leuprolide acetate for uterine fibroids. *N Engl J Med* 2012;366:421–32 [PEARL II study]
- <sup>2</sup> Baird D, Dunson D. Why is parity protective for uterine fibroids? *Epidemiology*. 2003;14(2):247-50.
- <sup>3</sup> Donnez J, Donnez O, Matule D, et al. Long-term medical management of uterine fibroids with ulipristal acetate. *Fertil Steril* 2015, Oct 23 [Epub ahead of print].
- <sup>4</sup> Courtoy G, Donnez J, Marbaix E, Dolmans MM. In vivo mechanisms of uterine myoma volume reduction with ulipristal acetate treatment. *Fertil Steril* 2015; **104**(2):426-34.
- <sup>5</sup> Yoshida S, Ohara N, Xu Q et al. Cell-type specific action actions of progesterone receptor modulators in the regulation of uterine leiomyoma growth. *Semin Reprod Med* 2010; 28(3):260-73.

